FERTILIZER TO-LOCATE-SOWING AND TRANSPLANTATION



FERTILIZER SOLUTION NP 5-30 with zinc (Zn)

DESCRIPTION

It is suggested the usage as fertirrigante in the first phase of the crop cycle or such in localization in the seeding and / or transplantation of the crops.

The product stimulates the development of radical apparatus, and applied in the phases of active growing favors the plastic formation of the plants, giving it strength and durability.

Phosphorus is a macroelement that intervenes in the process of lignification of the vegetative organs, ensuring the plant a greater resistance to adversity weather and to attacks by parts of the parasites.

The presence of polyphosphoric ions also allows to complexing microelements present in the soil, protecting by the phenomena of retrogradation and making them more easily available to crops.

METHOD 'AND DOSAGE

TREATMENTS RADICALS: in general in the uses of fertigation lend themselves effectively, at a dose of 55-120 kg / ha

Dosages for localized applications:

grain cereals (wheat, barley, oats) 55-110 kg / ha;

corn, sorghum 80-120 kg / ha; rice 80-120 kg / ha;

vegetables (tomatoes) 90-150 kg / ha

The doses given are the result of applied and calculated for distributions to normal volumes of water. For the correct application in the specific climate, soil and crop, it is advisable to consult your service technician. Respect the time of shortage.

COMPOSITION

Total Nitrogen (N)	5.0%
Ammonia Nitrogen (N)	3.0%
Urea Nitrogen (N)	2.0%
Phosphoric anhydride (P205) soluble in water	30.0%
Zinc (Zn) soluble in water	0.2%
Zinc (Zn) chelated by EDTA	0.2%



CE FERTILIZER







Chelating agent: EDTA Ph range which guarantees a good stability of the chelating fraction: 4-7.

The methods of analysis not reported are internal methods of the manufacturer.

Choncimer srl - F.ne Rocchetta, 63 - San Severino Marche (MC) tel. 0733636325 : e-mail:info@choncimer.it : P.IVA 01265640431